
HAIR: History of Animals using Isotope Records

(You are what you eat . . . plus a few ‰)

Thure CerlingUniversity of Utah





Thanks

- Linda Ayliffe, Gabe Bowen, Claire Cerling, Dylan Cerling, Lesley Chesson, Edna Ehleringer, Jim Ehleringer, Iain Douglas-Hamilton, Jackson King'oo, Passey, Dave Podlesak, Chris Remien, Jerry Schuster, Matt Sponheimer, Kevin Uno, George Wittemyer
- IsoForensics, Kenya Wildlife Service, National Science Foundation, Packard Foundation, Save-The-Elephants, Sheldrick Wildlife Trust

PNAS 10 Nov 2009

Cooperation and individuality among man-eating lions

Justin D. Yeaket^{1,1}, Broce D. Patterson², Keta Fox-Dobbs¹, Mercedes M. Okomura², Thure E. Cerling¹, Jonathan W. Moore¹, Paul L. Kochi², and Nathaniel J. Dominy^{1,1,1}

Toppertment of thickopy and Evolution fined (Museum or Repurs History, 146 Towns, Callle 1046, Taconia, Not Spirit IBC 1004, childed kimpsom, "Departic and Evolutionary Bridge, University and "Antiferoperage, University of Call

Refried by Krisson Howeley, Lebester,

Ecoperation is the conventions rious uses, a coalition of two aduficación, cooperatively killed dose "man-estaro of Tapon" have sino

MAN-EATERS
OF
TSAVO
Lt. Colonel J.H. Patterson, D.S.O.

Peter Capstick, Series Editor



sity of human prodution (up to 30% the of 1856) is site associated with , which may have further promoted nal prey under perturbed environ-

Sprittery les : Tarto : Tours

date that retails from high introspelus been reprinted in many minute to effects on the social behaviors of

compositive species are provely understand. Assemp lines, compositive behaviors are well known (4-7), yet the province of logganite (9), cheaters (9), and specialized hunters (7) within

1159 High Street, Santa Disk, CA 90094, "Department of Durings, efficient of Carrings, Entered of Space, Street, Stree

or JS, 3000 Devoted for ravies May 13, 3000.




PNAS 10 Nov 2009

Cooperation and individuality among man-eating lions

Justin D. Yeakel^{1,1}, Bruce D. Patterson², Kena Fox-Dobby¹, Mercedes M. Okomuru², Thure E. Cerling¹, Jonathan W. Moore¹, Paul L. Koch², and forthaniel J. Dominy^{1,1,1}

Toppertners of troology and Produtionary Storings, consensity of Carlifornia, Santa Cruz. 1759 High Street, Santa Cruz. CA 50004, "Organization of Country, Mind Shussays or September of September of September of September of August Storing, Color Storing, Storing Storing, Color Storing, Color Storing, Color Storing, Color Storing, Storing Storing, Storing Storing, Storing Storing, Storing Storing, Storing Stori

lighted by from transition, introducts of totals. Not calle title, 47, and approved beprovider 25, 2001 (increased for nation May 11, 2000)




"I'm trying to eat more vegetarians."


The New Yorker,14 Dec 2009


Elements are defined by the number of protons in the nucleus:


1=H, 2=He, 3=Li, 4=Be, 5=B, 6=C, 7=N, 8=O, and so on . . .





Isotopes are the different numbers of neutrons for a given proton configuration















ıllA

Boron

10.811



Isotope terminology

• $R_{phase} = {}^{13}C/{}^{12}C$ ratio in "phase"


• δ^{13} C (‰) = (R_{phase} / R_{standard} -1)*1000

• On Earth, 13 C ranges from 1.04% to about 1.14% (*ca.* δ^{13} C range from -65% to +25%)

Stable isotopes and animal behavior on different time scales

- A Traveler's Tale
- You are what you eat. Plus a few \%.
- Reaction progress variable
- Modern elephant isotope ecology
- NCIS forensics

δ^{13} C distinction between C₃ and C₄ plants





C₃ and C₄ photosynthesis

- C₃ plants
- most dicots
- cool-season grasses
- Foods: vegetables, fruits, beans, wheat, barley, rye, meat (from diet)



- rare in dicots
- tropical grasses and sedges
- Foods: maize, sorghum, sugar cane, millet, tef, fonio, and meat (from diet)









You are what you eat . . .

. . plus a few permil (%)



Ingredients:

Flour, Cornflour, Sugar, Hydrogenated Vegetable Oil, Ginger Powder, Salt, Invert Syrup, Lecithin, Chemicals, Synthetic Flavours & Permissible Food

Net Weight when packed 200 gms.

الكوثات دقيق، دقيق الذره اسكر، رستاق، محوق رنجيل املح اعصرا لستبن اكماشات روائح، الوان عداشه.

الوزن : ١٠٠٠ عندام

يرم كا آثار يمني كا آثار جيني ، يناسيتي ونك -- الأم

You

are.

Max. Retail Price

English Biscuit Manufacturers (Private) Limited Karachi - Pakistan.

MAX. RETAIL RS 7 35



New

Biscuits



Ingredients:

. . what you eat.

Flour, Cornflour, Sugar, Hydrogenated Vegetable Oil, Ginger Powder, Salt, Invert Syrup, Lecithin, Chemicals, Synthetic Flavours & Permissible Food Colours.

Net Weight when packed 200 gms.

What the traveler ate




What the traveler ate . . .





Movements between Mongolia and USA are recorded in beard hair












Univ Utah-BYU: controlled diets








Isotope enrichment ε*_{hair-diet} =3.1‰
















Isotope discrimination (air to plant) and enrichment (diet to animal tissue)



Controlled diet change UU – BYU

diet change of horses: C_3 to C_4 . Back to C_3 after 150 days. Ends with 7 day spike of C_4

• horse tail hair sectioned to 1 mm (ca. 1 day)

Team Utah-BYU

Dandy with Linda Ayliffe






Controlled diet change UU - BYU

- diet change: C_3 to C_4 . Back to C_3 after 150 days. Ends with 7 day spike of C_4
- horse tail hair sectioned to 1 mm (ca. 1 day)


Hair treated as $dN_i/dt = -\lambda_i N_i$

Isotope composition of hair predicted from known



-Ayliffe et al. 2004

Treat as "reaction progress": isotope values normalized from 0 to 1 (Ayliffe et al., 2004; Cerling et al 2007)



Model to directly calculate diet from isotope measurements of hair

Each pool changes as:


$$\delta_{i,(t)} = \delta_{i,(t-1)} e^{-\lambda_i(\Delta t)} + \delta_{D(t)} \left(1 - e^{-\lambda_i(\Delta t)}\right)$$

And thus diet is directly calculated as:

$$\delta_{\mathrm{D(t)}} = \frac{\left(\frac{\delta_{\mathrm{H}} + 1000}{\alpha_{\mathrm{HD}}} - 1000\right) - \left(\sum_{i=1}^{n} f_{i} \delta_{i, t-1} e^{-\lambda_{i} \Delta t}\right)}{\left(\sum_{i=1}^{n} f_{i} (1 - e^{-\lambda_{i} \Delta t})\right)}$$

Direct calculation of diet from sequence

- Horse moved from Brazil to Utah: July 2006
- Sampled April 2007
- Assume growth rate of 0.77 mm per day to match known time of movement
- Model of Cerling et al. (2007)





GPS collars and elephant hair










How do we get growth rates of hair?




Collect hair from same indviudal several times.

"Wiggle Match"


Can also correlate to other individuals QE is in same family unit as Anastasia (QE is mother of Anastasia)

Baton is passed to Cleopatra (sister)



Quantify the importance of grass during rainy season









Samburu

$$NDVI = \frac{(NRI-VIS)}{(NRI+VIS)}$$





Quantify the importance of grass during rainy season




Samburu


Data is for average ± 2 σ for 5 Samburu Reserve resident elephants









Crop raiding by Lewis occurs at night: outside the Imenti Forest




Crop raiding by Lewis occurs at night: outside the Imenti Forest






Diet, climate change, land-use change










Predicting behavior (diet)






Diet, climate change, land-use change







GPS tracking shows they left the park and were in community lands where they cannot compete with cattle: the grass is too short!



- Stable isotopes document the timing and progression of
- A. Rains
- B. Net Primary
 Productivity
- C. Water availab
- D. Diet respons
- E. Fertility



How do hair and ivory compare?





Other tissues? Compare hair and ivory.



- Amina: d. 27 Sept 2006
 - Tusk. 27 Sept 2006
 - Molar (M6). 27 Sept 2006
 - Hair. 27 Sept 2006; 16 June 2006; 1 Nov 2002;
 4 Feb 2002; 25 Jan 2001
 - GPS. 25 Jan 2001 to 27 Sept 2006 (intermittent)

Kevin Uno, UU



Misha: d. 9 Sept 2008

- Tusk. 9 Sept 2008
- Molar, M5 and M6
- Hair. None
- Travel. Moved from Vallejo CA to Salt Lake City UT 22 April 2005



Elephant ivory: longitudinal section

July 2004

milling striations


growth laminations

1000 microns

Jan 2005

Kevin Uno, UU

Amina: d. 27 Sept 2006




Kevin Uno, UU

How do you collect collect elephant hair?




5 tons vs. 1½ ton ("curb" weights)


"Rommel"



George Wittemyer's truck





What is response to change in location?




Change in location recorded in a single hair. (Beijing to Salt Lake City).

• Residents of different cities have characteristic isotope ratios in their hair








Tap water "isoscape" based on > 1000 municipal water supplies

USA Tap Water Map δ^{18} O, Bowen et al., 2007



Differences in the naturally occurring stable isotopes of water are "inherited" by local people in their hair

Tap water "isoscape" based on > 1000municipal water supplies

Dylan Cerling



Claire Cerling



Edna Ehleringer
























Differences in the naturally occurring stable isotopes of water are "inherited" by local people in their hair





"Saltaire Sally"



A local "cold case"

Hair, skull, some bones were found near Saltaire, Utah

Bundling of hair: increases resolution by increasing sample amount


Decreasing resolution by "averaging" samples with different growth rates and in different stages of growth (anagen versus catagen-telogen).

Growth rates (ca. 1 cm per month) in humans varies by ca. \pm 10% Approximately 10% of hairs are in "resting stage".

Model that follows:

90% anagen stage 60 ± 6 months

10% catagen-telogen 6 ± 1 months





-Menke, 2012





- Multiple hairs
- Gaussian distribution of growth rates and phases
- Signal increasingly "blurred" with increasing time

Hair Bundling (Remien thesis)











Signal recovered with inversion methods

but high frequency information can be lost

Example. high resolution record of elephant hair (ca. 3 years)

- A. Original
- B. Modeled: assume 10% growth rate variation
- C. Deconvolution using inverse methods






"Saltaire Sally"





A local "cold case"

Hair, skull, some bones were found near Saltaire, Utah





Original interpretation: Travel in N-S direction, about every 6 months






Original interpretation: Travel in N-S direction, about every 6 months






1. Inversion

2. Correct to "equilibrium"












You are what you eat (+ a few %)








Backup


NCIS (A Cold case, re-opened in 2008)




Hair from "Jane Doe", found in canal in Stanislaus County, CA on 11 September 1971 with 65 stab wounds.

Isotopes indicate limited travel in last year of life

Reconstructed travel history



Reconstructed travel history



MARY ALICE WILLEY and the BLACK LIBERATION ARMY


- Mary Alice Willey was born in Anaheim, CA in 1949; she moved to San Francisco, CA in **September 1969** to attend college. She grew disillusioned with school, dropped out, and became involved in the Black Power movement. In high school she had been known as Carol.
- 21 August 1971, she is thought to have smuggled a gun to George Jackson, a San Quentin prisoner, resulting in the killing of several prison guards and the death of Jackson during an escape attempt.
- 28 August 1971, a woman fitting the description of Mary Alice (but who gave the name Carol George) entered a San Francisco police station to report a missing purse. After completing paperwork, she signaled to individuals outside the station, who entered and killed policemen in apparent retaliation for George Jackson's death.
- 11 September 1971, the body of "Jane Doe" was discovered floating in a canal ditch in Stanislaus County.





San Francisco (Hetch-Hetchy

Reconstructed travel history



From hair

- last year of life
- *i.e.*, 1971

Who was this person?

From teeth

- ca. 6 years old
- *i.e.*, 1950


Anaheim, CA

MARY ALICE WILLEY and the BLACK LIBERATION ARMY


- Mary Alice Willey was born in Anaheim, CA in 1949; she moved to San Francisco, CA in September 1969 to attend college. She grew disillusioned with school, dropped out, and became involved in the Black Power movement. In high school she had been known as Carol.
- 21 August 1971, she is thought to have smuggled a gun to George Jackson, a San Quentin prisoner, resulting in the killing of coursel prison guards and the death of Jackson during an esca
- 28 August 1971, a woman fitting the description of M who gave the name Carol George) entered a San Franstation to report a missing purse. After completing pay signaled to individuals outside the station, who entere policemen in apparent retaliation for George Jackson's
- 11 September 1971, the body of "Jane Doe" was disin a canal ditch in Stanislaus County.
- Stable isotopes used for confirmation of DNA studies. ID established in 2009.


Prospects for reconstruction of diets using tooth enamel

Real input vs. modeled output



Real input vs. modeled output







Enamel does not form all at once

unit volume of enamel forms over time interval


Limit to sample resolution: can never get pure signal no matter what the sampling strategy

-Passey and Cerling, 2002





model for enamel maturation and sampling




Maturation parameters: Micro-Computerized Tomography


Micro-CT can be used to determine maturation parameters in developing tooth enamel

Corroborated by *in situ* density measurements


-Passey and Cerling, unpublished



Controlled diet change for rabbit. Forward model for teeth based on breath data and maturation parameters. Passey et al (2005).




Real input vs. modeled output



Inversion solution: (Meinke's "moving camera")

Set up input as matrix:

$$Am = d$$




The inverse solution to solve for the input is:

$$\mathbf{m}_{est} = \langle \mathbf{m} \rangle + \mathbf{A}^{T} \left[\mathbf{A} \mathbf{A}^{T} + \varepsilon^{2} \mathbf{I} \right]^{1} \left[\mathbf{d} - \mathbf{A} \langle \mathbf{m} \rangle \right]$$


Passey et al., 2005


Inversion method recovers original input



Laser ablation: micro-sampling


• Podlesak et al., 2007





Hippo tusk: Tsavo NP, Kenya


- 610 mm tusk
- Known facts: died during drought
- δ^{13} C shows gradual decline in diet quality, δ^{18} O shows gradual rise
- Reconstruction:
 - Severe diet stress
 - Severe water imbalance






Diet reconstruction

- 100 inversions: $\pm 1\sigma$
- lm = 45 to 65 mm
- 1a = 8 to 12 mm
- $x \pm 1 \text{ mm}$; $z\pm 0.2 \text{mm}$
- δ^{13} C: $\pm 0.1\%$ 0.









You are what you eat (+ a few ‰): you cannot escape your history



Modern: detailed history (to a single day)

- diet
- migration
- climate change
- land-use change

